

Background

Ash/Dust
Retrievals -
Puyehue
Eruption June
2011Comparisons
to MODIS
AODCross
SectionsCollocation
and Statistics

Dust contamination in AIRS v5 and v6 T(p) retrievals? Retrievals of Volcanic Ash - A work in progress

Eric S. Maddy^{1,2}, Sergio DeSouza-Machado³, Chris
Barnet², Larrabee Strow³, Scott Hannon³, Paul Schou³

¹STC, ²NOAA/NESDIS/STAR ³UMBC
email: eric.maddy@noaa.gov

November 14, 2012

Background and Motivation

Background

Ash/Dust
Retrievals -
Puyehue
Eruption June
2011

Comparisons
to MODIS
AOD

Cross
Sections

Collocation
and Statistics

- Sergio De-Souza Machado has shown that the AIRS V5 temperature and moisture profile products and surface temperature products have spurious biases relative to ECMWF when his dust score is large (qualitative assessment of the dust amount).
- We've extended results using N. Nalli's AEROSE radiosondes (Maddy, GRL, 2012) and have shown at previous meetings that the AIRS V5 temperature product is affected by dust. How does v6 behave in dusty situations? - Limited data, but I'll show results.
- Volcanic ash and Saharan dust spectral signatures are similar, so we would expect similar negative effects in the presence of volcanic ash.
- Can we simultaneously retrieval dust (or volcanic ash) and cloud using SARTAs scattering model? ... a work in progress.



Background

Ash/Dust
Retrievals -
Puyehue
Eruption June
2011

Comparisons
to MODIS
AOD

Cross
Sections

Collocation
and Statistics



E.Maddy_Fri Nov 9 10:05:42 EST 2012


Background

Ash/Dust
Retrievals -
Puyehue
Eruption June
2011

Comparisons
to MODIS
AOD

Cross
Sections

Collocation
and Statistics



E.Maddy_Fri Nov 9 10:05:42 EST 2012


Background

Ash/Dust
Retrievals -
Puyehue
Eruption June
2011

Comparisons
to MODIS
AOD

Cross
Sections

Collocation
and Statistics



E.Maddy: Fri Nov 9 10:05:51 EST 2012


Background

Ash/Dust
Retrievals -
Puyehue
Eruption June
2011

Comparisons
to MODIS
AOD

Cross
Sections

Collocation
and Statistics



E.Maddy: Fri Nov 9 10:05:52 EST 2012


Background

Ash/Dust
Retrievals -
Puyehue
Eruption June
2011


Comparisons
to MODIS
AOD

Cross
Sections

Collocation
and Statistics



2012/09/20 vs. ECMWF (dusty cases)



EMaddy Fri Nov 9 10:06:12 EST 2012



Background

Ash/Dust
Retrievals -
Puyehue
Eruption June
2011

Comparisons
to MODIS
AOD

Cross
Sections

Collocation
and Statistics



EMaddy_Fri Nov 9 10:06:13 EST 2012



Background

Ash/Dust
Retrievals -
Puyehue
Eruption June
2011

Comparisons
to MODIS
AOD

Cross
Sections

Collocation
and Statistics



EMaddy Fri Nov 9 10:06:22 EST 2012



Background

Ash/Dust
Retrievals -
Puyehue
Eruption June
2011

Comparisons
to MODIS
AOD

Cross
Sections

Collocation
and Statistics



EMaddy Fri Nov 9 10:06:23 EST 2012

Background

Ash/Dust
Retrievals -
Puyehue
Eruption June
2011Comparisons
to MODIS
AODCross
SectionsCollocation
and Statistics

Retrieval of dust/ash from AIRS

- Use an unconstrained (or loosely constrained) optimal estimation algorithm to retrieval dust (ash) particle size, loading and height.
- SARTA's scattering algorithm currently in use.
- Also can estimate ice/cirrus in addition to dust - now using some brute-force type retrievals from UMBC (Sergio).

An example of retrievals from Puyehue - dust heights


Background

Ash/Dust
Retrievals -
Puyehue
Eruption June
2011

Comparisons
to MODIS
AOD

Cross
Sections

Collocation
and Statistics



Background

Ash/Dust
Retrievals -
Puyehue
Eruption June
2011Comparisons
to MODIS
AOD

Discussion

Cross
SectionsCollocation
and Statistics

- Select all MODIS pixels falling within 15 kilometers of AIRS footprint FOV.
- Compare the individual MODIS pixels MYD04 product for Aerosol OD comparisons to our volcanic ash loading (Either Deep_Blue_Aerosol_Optical_Depth_550_Land or Effective_Optical_Depth_Average_Ocean).



Background

Ash/Dust
Retrievals -
Puyehue
Eruption June
2011Comparisons
to MODIS
AOD

Discussion

Cross
SectionsCollocation
and Statistics

Ash loadings - Comparison to MODIS



Background

Ash/Dust
Retrievals -
Puyehue
Eruption June
2011Comparisons
to MODIS
AOD
DiscussionCross
SectionsCollocation
and Statistics

Discussion and Future Work

- AIRS v5 is severely affected by dust - most of the errors come from the first guess PC regression.
- AIRS v6 is moderately affected by dust. Appears that the SCCNN is somewhat immune to dust contamination - AIRS v6 physical retrieval has slightly larger RMS uncertainties as compared to the SCCNN.
- We suggest flagging dusty cases in AIRS L2.
- Work in progress, but we have shown AIRS can retrieve dust/ash loadings that are comparable to MODIS AOD.
- More work is needed to optimize the algorithm/constraints - but simultaneous dust + L2 retrievals should enable more accurate retrievals in the presence of dust.

AIRS DUST

E. Maddy et
al.

Background

Ash/Dust
Retrievals -
Puyehue
Eruption June
2011

Comparisons
to MODIS
AOD

Discussion

Cross
Sections

Collocation
and Statistics

Discussion and Future Work


Background

Ash/Dust Retrievals - Puyehue Eruption June 2011

Comparisons to MODIS AOD

Cross Sections

Collocation and Statistics




Background

Ash/Dust
Retrievals -
Puyehue
Eruption June
2011

Comparisons
to MODIS
AOD

Cross
Sections

Collocation
and Statistics



Background

Ash/Dust
Retrievals -
Puyehue
Eruption June
2011Comparisons
to MODIS
AODCross
SectionsCollocation
and Statistics
AEROSE


Collocation Strategy

- We select all retrievals within $\delta t = \pm 3$ hr, $\delta r = 150$ km of a Microtops Sunphotometer measurement and corresponding collocated RAOB.
- We calculate mean bias tendency and standard deviation statistics for data that fall within 4 AOD bins.
- We also use the RAOB profiles + ECMWF surface temperature and compare cloud-cleared radiances from AIRS and IASI to the forward calculations for each of the 4 AOD bins

Background

Ash/Dust
Retrievals -
Puyehue
Eruption June
2011Comparisons
to MODIS
AODCross
SectionsCollocation
and Statistics


AEROSE



Background

Ash/Dust
Retrievals -
Puyehue
Eruption June
2011Comparisons
to MODIS
AODCross
SectionsCollocation
and Statistics


AEROSE



Background

Ash/Dust
Retrievals -
Puyehue
Eruption June
2011Comparisons
to MODIS
AODCross
SectionsCollocation
and Statistics


AEROSE



Background

Ash/Dust
Retrievals -
Puyehue
Eruption June
2011Comparisons
to MODIS
AODCross
SectionsCollocation
and Statistics

AEROSE



Ship Tracks

Background


Ash/Dust Retrievals - Puyehue Eruption June 2011

Comparisons to MODIS AOD

Cross Sections

Collocation and Statistics

AEROSE



Correlative Data of Interest

Dedicated RAOBs

- Typically 4/day at 01:30, 09:30, 13:30, 21:30
- Over 250 PTU soundings in 2009, 2010, 2011
- not assimilated, decoupled from land-based RAOBs - thus truly independent

Ozone sondes

- Over 64 soundings in 2009, 2010, and 2011
- 113 O₃ soundings to date

Microtops Sunphotometer

- Multi-channel aerosol optical depth (AOD)
- NASA/GSFC AERONET Maritime Aerosol Network (MAN) methodology and QC applied to retrieve a standardized AOD.